

9400265

TO ALL TO WHOM THESE PRESENTS; SHALL COME;

USDA-NRCS Aberdeen Plant Materials Center

MICTORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A part hereof, and the various requirements of LAW in such cases made and provided have been COMPLIED WITH, AND THE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT variety protection under the LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC replenishment of viable basic seed of the variety in a public repository as provided by LAW, the SHILLO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR TING II, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE URPOSE, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT IN THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) D BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE ERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321

WHEATGRASS, INTERMEDIATE

'Rush'

In Testimony Therest, I have hereunto set my hand and caused the seal of the Hant Inviety Archivetion Office to be affixed at the City of Washington, D.C. this twenty-fourth day of April, in the year of our Lord two thousand one.

ET SEQ.

Plant Varioty Protection Office Agricultural Markoting Service

AODITCE LOCALLY. Include form number and	adition date on all	reproductions.	OMB APPROVED NO. 0581-005	
U.S. DEPARTMENT AGRICULTURAL M. SCIENCE	Application is required in order to determine it a plant variety protection certificate is to be issued (7 U.S.C.			
APPLICATION FOR PLANT VAI	2421). Information is held confidential until certificate is Issued (7 U.S.C. 2426).			
1. NAME OF APPLICANT(S) (as it is to appear on the Certificat	e)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NO.	3. VARIETY NAME	
USDA-SCS Aberdeen Plant Materials	S Center	PI 575702	Rush	
4. ADDRESS (street and no. or R.F.O. no., city, state, and ZIP)	***************************************	5. PHONE (include area code)		
P.O. Box 296		(208) 397-4133	PVPO NUMBER	
Aberdeen, ID 83210-0296			9400265	
			Date Que 1, 1994	
6. GENUS AND SPECIES NAME	7. FAHILY NAME (B	otanical)	N G G A.M. D P.M.	
Elytrigia intermedia	Poaceae		F Filing and Examination Fee:	
8. CROP KIND NAME (Common Name)		9. DATE OF DETERMINATION		
Intermediate Wheatgrass		4/90	S Date	
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FOR	LL OF ORGANIZATION	1	- aug. 1, 1994	
association, etc.)		Trooperation, partitionally,	Certificate Fee:	
U.S. Government			½	
11. IF INCORPORATED, GIVE STATE OF INCORPORATION		12. DATE OF INCORPORATION	D Date	
13. NAME AND ADORESS OF APPLICANT REPRESENTATIVE(S	IE ANY TO SERVE	N THIS ABOUT AND BECSIVE	E ALL DADEDS	
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUB a	ity wnership a Seed Sample mailed	to Plant Variety Protection Office	(206) 397-4133	
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VA Plant Variety Protection Act) YES (# "YES," answer	CRIETY BE SOLD BY V	ARIETY NAME ONLY AS A CLASS (
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE			S OF PRODUCTION BEYOND BREEDER SEED?	
LIMITED AS TO NUMBER OF GENERATIONS?			GISTERED 🛣 CERTIFIED	
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION TYPES (II "YES," through Plant Variety Protection NO		N THE U.S.? Int Act. Give date:		
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR YES (If "YES," GIVE NAMES OF COUNTRIES AND DA	SALE, OR MARKETE	D IN THE U.S. OR OTHER COUNTR	NES?	
□ NO				
 The applicant(s) declare(s) that a viable sample of basic seed such regulations as may be applicable. 	s of this variety will be	furnished with the application and t	will be replenished upon request in accordance with	
The undersigned applicant(s) is (are) the owner(s) of this sexu in section 41, and is entitled to protection under the provision	ally reproduced novel p a of section 42 of the	plant variety, and believe(s) that the Plant Variety Protection Act.,	variety is distinct, uniform, and stable as required	
Applicant(s) is-(are) informed that false representation herein	can jeopardize protect	ion and result in penalties.	·	
IGNATURE OF APPLICANT (OWNer(s))		CAPACITY OR TITLE	DATE	
Coul-fl aluster		_4	6/16/94	
GNATURE OF APPLICANT [Owings]		State Conservationia CAPACITY OR TITLE		
OF AFFEIGANT [OWING(S)]		WACIT ON IILE	DATE	
470 (06.93) Editions of Forms S-470 (3/86) and CSSD	470 (5:80) are to b	o dostava d		

A. ORIGIN AND BREEDING HISTORY (Revised)

'Rush' (PI 575702) Elytrigia intermedia (Host) Nevski, was selected from very broad germplasm (PI 281863) which has been in the US since 1962. This germplasm was entered into the PI system as Agropyron junceum (L.) Beauv. The source was the German Botanical Garden, Berlin.

Through several cycles, Rush was selected for numerous, broad leaves, superior seedling emergence, (emerging 3 to 5 days earlier and in greater numbers), uniform performance and consistent rhizomatous form compared to the original broad-based PI 281863 germplasm. Final selection cycles for Breeder material were made vegetatively.

Rush has been observed to be stable and uniform over five generations since 1988. The variants which have been found in Rush include glaucous leaves (approximately 5 percent) and non-rhizomatous plants (less than 5 percent). Under irrigated conditions approximately 30 percent of Rush plants exhibit wide basal leaves ranging from 17 - 20 mm.

B. NOVELTY STATEMENT for 'Rush' Intermediate Wheatgrass (Revised)

The most similar existing variety to Rush intermediate wheatgrass is 'Oahe'. Oahe was selected at South Dakota Agricultural Experiment Station, Brookings and was released in 1961. Oahe was selected from a Russian introduction identified as PI 98568. Oahe, a four-clone synthetic, was developed from two cycles of selection for high seedset, forage, yield, and rust resistance within a population of high-seed-producing plants. It is described as being uniformly blue green, vigorous, rhizomatous, drought tolerant, winter hardy, and produces high yields of large seed (Grass Varieties in the United States, Agriculture Handbook No. 170, Soil Conservation Service, revised November, 1994).

Table 1 is a comparison of morphological characteristics of Rush and Oahe intermediate wheatgrass. Characteristics were measured during the 1996 growing season at two separate locations, The Curlew Test Site and the Aberdeen Plant Materials Center Display Nursery, both located in southeastern Idaho.

Rush is similar to Oahe in terms of stem diameter, flag leaf width and length, leaf width below flag leaf, stem exsertion and flowering date. Rush and Oahe are also similar in color (5GY/5, Munsell Color Chart for Plant Tissues), however Oahe is uniformly glaucous while Rush exhibits the glaucous characteristic in approximately five percent of the plants. Rush is also similar to Oahe in terms of drought tolerance, vigor and winter hardiness.

The differentiating characters between Rush and Oahe are basal diameter, plant height and seed weight. Rush produces a significantly larger basal diameter as compared to Oahe and Rush is a significantly taller plant than Oahe. These characteristics were significantly different at both locations. Rush also has larger seed than Oahe as shown on Table 1.

Table 1. Morphological Comparison of 'Rush' and 'Oahe' Intermediate Wheatgrass

CURLEW TEST SITE 11

PMC DISPLAY NURSERY 2/

	RUSH	OAHE	COMPUTED ³ ∕		RUSH	OAHE	COMPUTED 3/
,			T _(0.01)				T _(0.01)
x =	10.4	6.8	5.83	χ=	12.5	9.4	6.01
s=	2.5	2.2		s=	1.7	2.3	
-χ̈=	61.9	49.3	5.66	Σ =	105.3	94.8	3.34
S=	9.5	7.6		s=	9.2	14.4	
<u>x</u> =	2.8	2.5	2.18	x =	2.7	3.0	1.49
s≖	0.6	0.6		s=	0.7	0.7	
- =	5.8	5.5	0.84	x =	7.0	8.5	2.81
S=	1.5	1.2		S=	1.9	2.2	
x =	16.0	13.0	2.49	z =	14.3	13.7	0.43
s=	5.4	3.8		s=	5.0	5.7	
χ =	5.6	5.2	0.96	x =	5.4	5.6	0.47
s=	1.8	1.7		s=	1.9	1.3	
x =	5.1	4.8	1.27	x =	5.4	6.2	3.53
\$ =	1.0	1.2		s=	0.7	0.9	
<u></u> =	15.7	14.0	1.93	x =	20.1	23.6	3.24
S =	3.5	3.3		S=	3.0	5.0	
<u></u> =	16.0	9.7	2.67	Σ=	35.5	28.2	2.03
S=	9.8	8.4		s=	14.7	13.3	
x =	12.7	11.4	0.92	र्⊼=	39.6	40.7	0.66
s=	5.7	5.2		s=	5.4	7.4	
	5GY/5	5GY/5			5GY/5 to	5GY/5	(Munsell Color
	Approx. 5%	100%			7.5 GY/5		Chart for Plant
	glaucous	glaucous			approx. 5%	100%	Tissues).
					glaucous	glaucous	
		<u></u> .					
	Jul-10	Jul-10			Jul-1	Jul-1	
					6.87	5.01	
	S= X= S= X= S= X= S= X= X= <td>X= 10.4 s= 2.5 X= 61.9 s= 9.5 X= 2.8 s= 0.6 X= 5.8 s= 1.5 X= 16.0 s= 1.8 X= 5.1 s= 1.0 X= 15.7 s= 3.5 X= 16.0 s= 9.8 X= 12.7 s= 5.7 5GY/5 Approx. 5% glaucous</td> <td>X= 10.4 6.8 s= 2.5 2.2 X= 61.9 49.3 s= 9.5 7.6 X= 2.8 2.5 s= 0.6 0.6 X= 5.8 5.5 s= 1.5 1.2 X= 16.0 13.0 s= 5.4 3.8 X= 5.6 5.2 s= 1.8 1.7 X= 5.1 4.8 s= 1.0 1.2 X= 15.7 14.0 s= 3.5 3.3 X= 16.0 9.7 s= 9.8 8.4 X= 12.7 11.4 s= 5.7 5.2 SGY/5 Approx. 5% glaucous glaucous glaucous</td> <td>$\begin{array}{c ccccccccccccccccccccccccccccccccccc$</td> <td>$\begin{array}{c ccccccccccccccccccccccccccccccccccc$</td> <td>$\begin{array}{c ccccccccccccccccccccccccccccccccccc$</td> <td>$\begin{array}{c ccccccccccccccccccccccccccccccccccc$</td>	X= 10.4 s= 2.5 X= 61.9 s= 9.5 X= 2.8 s= 0.6 X= 5.8 s= 1.5 X= 16.0 s= 1.8 X= 5.1 s= 1.0 X= 15.7 s= 3.5 X= 16.0 s= 9.8 X= 12.7 s= 5.7 5GY/5 Approx. 5% glaucous	X= 10.4 6.8 s= 2.5 2.2 X= 61.9 49.3 s= 9.5 7.6 X= 2.8 2.5 s= 0.6 0.6 X= 5.8 5.5 s= 1.5 1.2 X= 16.0 13.0 s= 5.4 3.8 X= 5.6 5.2 s= 1.8 1.7 X= 5.1 4.8 s= 1.0 1.2 X= 15.7 14.0 s= 3.5 3.3 X= 16.0 9.7 s= 9.8 8.4 X= 12.7 11.4 s= 5.7 5.2 SGY/5 Approx. 5% glaucous glaucous glaucous	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

The Curlew test site is a dryland plant testing site located approximately 70 miles southeast of Aberdeen, Idaho. Average annual precipitation is 12-16 inches.

The PMC Display Nursery is located at the Aberdeen, Idaho Plant Materials Center Farm. These plots are irrigated to achieve the equivilent of 12-16 inches annual precipitation.

The T-test was used to compare Rush and Oahe. Thirty individual plants from each cultivar were evaluated at each testing site during the 1996 growing season. Degrees of Freedom (d.f.) = 58, level of significance = 0.01. Tabular $T_{(0.01)} = 2.66$. Computed T values in **bold** indicate plant characters which are significantly different between Rush and Oahe.

 $[\]frac{4\ell}{2}$ 1000 seed weight data from Natural Resources Conservation Service Technical Guide, section IV Standards and Specifications.

C. OBJECTIVE DESCRIPTION OF VARIETY

'RUSH' INTERMEDIATE WHEATGRASS (ELYTRIGIA INTERMEDIA (Host) Nevski

Name of Applicant: USDA-Soil Conservation Service

Temporary Designation: NA

Plant Introduction Number: PI-575702

Variety Name: Rush

- 1. SPECIES: Elytrigia intermedia, synonyms include Thinopyrum intermedium, and Elymus hispidus. Former name, no longer accepted, is Agropyron intermedium.
- 2. CYTOLOGY: Allohexaploid, 2n=42, S and X genomes for Triticeae tribe of the Poaceae family.
- 3. ADAPTATION: USDA-SCS Major Land Resource Areas; A3, A5, B6, B8-10, B13, D21, D22, D25, D28a, and E43-51, where rainfall exceeds 12 inches. Under irrigation, it will grow in all A, B, D, and E regions. It is probably adapted to many areas across the northern tier of states and southern Canada.
- 4. MATURITY: Anthesis is approximately the same as other intermediate wheatgrasses in southern Idaho. Seed maturity is 5-10 days later than Tegmar, Greenar, and Greenleaf.
- 5. PLANT HEIGHT: Average height of mature culm: 140 cm. Range 110-170 cm. Approximately 10 cm taller than Greenar or Greenleaf
- 6. GROWTH HABIT: Erect, densely tillered, leaves basal and on culms, somewhat lax.
- 7. RHIZOMES: Moderately rhizomatous, 1-3 dm per year.
- 8. LEAF BLADE:

Color: Blue-green

Glaucous: Yes, variable

Glabrous: No

Length: 30 - 40 cm

Width: 10 - 17 mm

9. LEAF SHEATH:

Auricles: Very small, 1.0 mm

Ligule: Short, 0.5 - 1.0 mm, square, minutely toothed

- 10. INFLORESCENSE: Spike, narrow and erect, spikelets diverging slightly at maturity, 15 -25 cm long
- 11. LEMMA:

Length: 10 - 12 mm

√ Width: 32 mm

Awn length: 1 - 2 mm

12. SEED: 7.5 g./ 1000 seeds

E. STATEMENT OF THE BASIS OF THE APPLICANTS OWNERSHIP

The variety for which PVP is being requested was developed by the staff of the U.S. Dept. of Agriculture, Soil Conservation Service, Plant Materials Center in Aberdeen Idaho. The leaders in this development were Gary Young, PMC Mgr, J. Chris Hoag, and Loren St John, PMC Asst Mgrs. As US government employees, these persons retain no ownership rights to this variety. All rights to this variety are held by and remain with the U.S. Dept. of Agriculture, Soil Conservation Service.

